

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: <http://www.elsevier.com/locate/euprot>

## Mini-review

# Biomarker discovery in asthma and COPD: Application of proteomics techniques in human and mice



Steven Haenen<sup>a,b</sup>, Elke Clynen<sup>c</sup>, Benoit Nemery<sup>b</sup>, Peter H.M. Hoet<sup>b</sup>, Jeroen A.J. Vanoirbeek<sup>b,\*</sup>

<sup>a</sup> KU Leuven, Laboratory of Drug Disposition & Delivery, Leuven, Belgium

<sup>b</sup> KU Leuven, Centre for Environment and Health, Leuven, Belgium

<sup>c</sup> Hasselt University, Biomedical Research Institute (BIOMED), Hasselt, Belgium

## ARTICLE INFO

## Article history:

Received 23 December 2013

Received in revised form

22 April 2014

Accepted 28 April 2014

Available online 5 June 2014

## Keywords:

Proteomics

Asthma

COPD

Human

Animal models

## ABSTRACT

The use of advanced proteomics approaches in the search for biomarkers in chronic lung diseases, such as asthma and COPD, is rather limited. Asthma and COPD are complex disorders, which can be subdivided into several phenotypes. This results in a heterogeneity of differentially expressed biological molecules. Furthermore, genetic differences between animals and humans make 'translation' of possible biomarkers challenging. Yet, the improved sensitivity and high throughput of proteomic techniques could be an important asset for (new) protein biomarker discovery in either human or animal models. We have reviewed the literature that reported the use of different proteomics approaches performed on samples obtained from humans and murine models in asthma and COPD research for the discovery of new biomarkers of diseases, biomarkers of sensitization or for the refinement of treatment. There is an increasing trend in the use of proteomics to explore new biomarkers of asthma or COPD. Although several murine models have been developed to study these lung diseases, and proteomics studies have been performed, 'translation' of identified candidate biomarkers into clinical studies is often lacking.

© 2014 The Authors. Published by Elsevier B.V. on behalf of European Proteomics Association (EuPA). This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

## Contents

1. Introduction .....	102
2. Proteomics approaches in asthma and COPD .....	102
3. Methodology .....	107

\* Corresponding author at: KU Leuven, Department of Public Health and Primary Care, Centre for Environment and Health, Herestraat 49 Mailbox 706, B-3000 Leuven, Belgium. Tel.: +32 16 33 01 96; fax: +32 16 33 08 06.

E-mail address: [jeroen.vanoirbeek@kuleuven.be](mailto:jeroen.vanoirbeek@kuleuven.be) (J.A.J. Vanoirbeek).

<http://dx.doi.org/10.1016/j.euprot.2014.04.008>

2212-9685/© 2014 The Authors. Published by Elsevier B.V. on behalf of European Proteomics Association (EuPA). This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

4. Mice vs. men .....	108
5. Sample origin .....	108
6. Interpretation of results .....	109
7. Future .....	109
8. Conclusion .....	109
Conflict of interest statement .....	110
Transparency document .....	110
References .....	110

## 1. Introduction

A rapid increase in the use of proteomics in different research domains such as cancer, invertebrate immunology and plant physiology was made possible thanks to the completion of several genomes, technical advances in instrumentation (increased sensitivity) and methods (purification, separation, identification) and the development of new bioinformatical tools [1–5]. In 2013, almost 5000 papers were published using a proteomics technique (*Pubmed search: Publication date: 2013; Keyword: Proteomics*). Yet, the use of advanced proteomics approaches in the search for biomarkers in chronic lung diseases, such as asthma and COPD, is rather limited. This limitation is attributable to multiple factors. First, asthma and COPD are complex disorders which are subdivided into several phenotypes [6–8]. This heterogeneity results in differential expression of biological effector molecules, namely proteins and peptides. Moreover, human samples are inherently characterized by a larger heterogeneity as compared to murine samples due to a different genetic background. Finally, ethically and practically, it is not feasible to obtain large numbers of human samples hampering large proteomic studies [9].

Nevertheless, the sensitivity and high throughput of proteomics techniques could be useful to discover (new) biomarker proteins in humans or animal models, previously not associated with a disease phenotype, at a time point much earlier than current existing diagnostic tools can do [10]. Despite efforts to develop animal models to model different human diseases, translation of (proteomics) research evidence from animals to humans is difficult and time consuming. Only one out of three animal studies evaluating medical interventions for human diseases have been replicated in clinical trials with an average time span of seven years to complete the translation from animals to humans [11].

We have reviewed the literature that reported the use of different proteomics approaches performed on samples obtained from humans and murine models in asthma and COPD research for the discovery of new biomarkers of diseases, biomarkers of sensitization or for the refinement of treatment.

Asthma affects more than 300 million people worldwide, whereas COPD is predicted to be the third most leading cause of death by 2030 (WHO; [www.who.int](http://www.who.int)).

Asthma, characterized by a reversible airway obstruction, a non-specific airway hyperreactivity and an airway inflammation, is orchestrated by the interplay of numerous cells, from both the innate immune system (e.g. dendritic cells, macrophages, mast cells, eosinophils and neutrophils) and

the adaptive immune system (T cells and B cells) [12–14]. Clinical manifestations of asthma are episodic wheezing, coughing, shortness of breath and non-specific hyperreactivity to e.g. cold air or cigarette smoke. Asthma can be subdivided in different classes (allergic asthma, non-allergic asthma, occupational asthma, etc.). Although different phenotypes of asthma have already been distinguished, the exact molecular mechanisms underlying the disease remain unclear [14].

Chronic obstructive pulmonary disease (COPD) is characterized by a progressive airflow limitation that is partially reversible, a chronic lung inflammation and systemic effects [6]. Although the main risk factor to develop COPD is smoking tobacco, with more than 90% of COPD patients having a smoking history, other factors such as maternal smoking, childhood asthma and outdoor air pollution have recently been associated with the development of COPD [15].

The prevalence of both asthma and COPD is increased during the last decades, indicating that current state-of-the-art techniques for diagnosis and management are suboptimal [16]. Current diagnostic tools such as spirometry fail to pinpoint the early onset of chronic lung diseases resulting in late diagnosis when adverse clinical symptoms have already been established. Moreover, the few existing molecular markers lack in sensitivity and specificity (e.g. nitric oxide, eosinophil cationic protein and eosinophil peroxidase) [16,17].

## 2. Proteomics approaches in asthma and COPD

Clinically, it has been estimated that almost 10% of the genome is directly involved in disease pathogenesis. Therefore, detecting distinct protein biomarkers, encoded by the genome, in certain pathologies can contribute to disease detection, monitoring disease progression and response to treatment [18].

So far, most proteomic studies performed in either patient cohorts or murine models of asthma and COPD have investigated established disease (Tables 1 and 2). Several of the differentially expressed proteins in these studies, could be linked with hallmarks of allergic asthma such as airway hyperreactivity, lung inflammation [19–21] and remodeling [22–24]. Proteomic changes due to long term cigarette smoke exposure in murine models could be related to lung damage and smoke-induced lung irritation [25,26]. Proteomic screening in humans suffering from COPD most often focused on samples obtained from the respiratory system, such as bronchoalveolar lavage (BAL), lung tissue or cells, epithelial lining fluid or induced sputum. Some studies have also analyzed potential

**Table 1 – Overview of proteomics studies in animal models of asthma and COPD.**

Reference	Species	Disease	Inducer	Intervention	Sample	Technique	Main findings	Up/down <sup>a</sup>
Houtman et al. [42]	Mouse	Non-allergic asthma	DNFB		Lung	2D-GE Silver staining	Coronin 1A Vinculin Gelsolin	↑ ↑ ↓
Roh et al. [43]	Mouse	Allergic asthma	OVA	DMS	Lung	2D-GE Silver staining	T complex polypeptide 1 EH-domain containing protein 4 Plasminogen	↑ ↓ ↓
Signor et al. [19]	Rat	Allergic asthma	OVA	Endotoxin	BAL	2D-GE Coomassie staining	Fetuin A Fetuin B Haptoglobin	↑ ↑ ↑
Jeong et al. [45]	Mouse	Allergic asthma	OVA		Lung	2D-GE Coomassie staining	Clara cell 10 kDa secretory protein Chitinase 3-like 3 Chitinase 3-like 4	↓ ↑ ↑
Zhao et al. [60]	Mouse	Allergic asthma	OVA		BAL	2D-GE Silver staining	Lungkine Chitinase 3-like 3 Chitinase 3-like 4 Calcium-activated chloride channel regulator 1	↑ ↑ ↑ ↑
Greenlee et al. [50]	Mouse MMP2 <sup>-/-</sup> MMP9 <sup>-/-</sup>	Allergic asthma	OVA		BAL	2D-DIGE	Chitinase 3-like 3 Calgranulin A	↑ ↑
Zhao et al. [36]	Mouse	Allergic asthma	OVA	DMS	BAL	2D-GE Silver staining	Chitinase 3-like 3 Chitinase 3-like 4 Vitamin D binding protein	↓ ↓ ↑
Liu et al. [61]	Mouse	Allergic asthma	OVA	Salbutamol	Lung	2D-GE Silver staining	Hemopexin Rho GDP-dissociation inhibitor 2 Peroxiredoxin 5	↓ ↑ ↑
Zhang et al. [25]	Rat	COPD	Cigarette smoke		Lung	2D-GE Coomassie staining	Thioredoxin Peroxiredoxin 6 α-Enolase	↑ ↑ ↓
Calvo et al. [23]	Mouse	Allergic asthma	OVA		Lung	SELDI-TOF MS	Found inflammatory zone 1 Calcyclin Clara cell 10 kDa secretory protein	↑ ↑ ↑

Table 1 (Continued)

Reference	Species	Disease	Inducer	Intervention	Sample	Technique	Main findings	Up/down <sup>a</sup>	
Xu et al. [26]	Rat	COPD	Cigarette smoke		Lung	2D-GE	RAGE	↑	
			Radon			Coomassie staining	Thioredoxin	↑	
							Calcyclin	↑	
Haenen et al. [20]	Mouse	Occupational asthma	TDI		Aur LN	2D-DIGE	Lymphocyte specific protein 1	↓	
					BAL		Vitamin D binding protein	↑	
					Serum		Hemopexin	↑	
Calvo et al. [24]	Mouse	Allergic asthma	OVA		Lung	2D-DIGE	78 kDa glucose-regulated protein precursor	↑	
Tewari et al. [35]	Mouse	COPD	Cigarette smoke		Plasma	2D-DIGE	Fibrinogen	↓	
							α-1-antitrypsin	↑	
							Arginase	↓	
Louten et al. [21]	Mouse	TSLP Tg mice	–		BAL	1D-GE	Chitinase 3-like 3	↑	
							Chitinase 3-like 4	↑	
	Macaque		HDM		BAL	LC-MS/MS	Cyclophilin A	↑	
							Cofilin A	↑	
Haenen et al. [41]	Mouse	Occupational asthma	TDI		Aur LN	2D-DIGE		1 sens	2 sens
					Serum		Coronin 1a	↑	↑
							Lymphocyte specific protein 1	–	↓
							Hemopexin	↑	↓

<sup>a</sup> Differences presented are always comparisons from a certain disease vs. healthy (e.g. asthma vs. control).

Identification of differentially expressed proteins was performed via mass spectrometry (MALDI-TOF MS or tandem mass spectrometric analysis).

Abbreviations: 1D-GE, one-dimensional gel electrophoresis; 2D-GE, two-dimensional gel electrophoresis; 2D-DIGE, two-dimensional difference gel electrophoresis; Aur LN, auricular lymph nodes; BAL, bronchoalveolar lavage; COPD, chronic obstructive pulmonary disease; DMS, dexamethasone; DNFB, dinitrofluorobenzene; EH-domain, epidermal growth factor receptor substrate 15; HDM, House Dust Mite; LC-MS/MS, liquid chromatography mass spectrometry; MMP, matrix metalloproteinase; OVA, ovalbumin; RAGE, receptor for advanced glycation endpoints; SELDI-TOF MS, surface enhanced laser desorption/ionization – time of flight mass spectrometry; sens, sensitization; TDI, toluene-2,4-diisocyanate; TSLP Tg, thymic stromal lymphopoietin transgenic.

**Table 2 – Overview of proteomics studies in human asthma and COPD.**

Reference	Disease	Sample	Technique	Main findings	Up/down <sup>a</sup>
Merkel et al. [28]	COPD	BAL	SELDI-TOF MS	Neutrophil defensin 1 Neutrophil defensin 2 Calgranulin A Calgranulin B Clara cell phospholipid-binding protein	↑ ↑ ↑ ↑ ↓
Candiano et al. [44]	Allergic asthma	Bronchial epithelial cells ( <i>IL-4 stimulated</i> ) BAL	2D-GE Coomassie staining	Gelsolin (45 kDa) Gelsolin (85 kDa) Gelsolin (45 kDa) after allergen challenge Gelsolin (85 kDa) after allergen challenge	↑ ↑ = ↓
Steiling et al. [29]	COPD	Bronchial epithelial cells	1D-GE LC-MS/MS	PLUNC Clara cell 10 kDa secretory protein Prolyl 4-hydroxylase beta subunit	↓ ↓ ↑
Larsen et al. [62]	Mild asthma	BAL	2D-GE Coomassie or silver staining	Haptoglobin	↑
Larsen et al. [46]	Mild asthma Scleroderma (SSc)	Fibroblasts cultured in BAL & bronchial biopsies	2D-GE Coomassie or silver staining	Alpha-smooth muscle actin	↑
Jeong et al. [47]	Allergic asthma	Peripheral T lymphocytes	2D-GE Silver staining	Phosphodiesterase 4 Thioredoxin 2 Glutathione S transferase M3	↑ ↑ ↓
Ohlmeier et al. [48]	COPD	BAL	2D-GE Silver staining	Surfactant protein A	↑
Gray et al. [27]	Allergic asthma COPD CF	Sputum	SELDI-TOF MS	Calgranulin A, B & C Clara cell 10 kDa secretory protein Proline rich salivary peptide	↑ ↓ ↓
Wu et al. [22]	Allergic asthma	BAL after segmental airway challenge	LC-MS/MS	Matrix metalloproteinase-9 Calgranulin B Tissue inhibitor of metalloproteinases 1	↑ ↑ ↑
Bandow et al. [63]	COPD	Plasma	2D-GE Sypro Ruby	Plasma retinal-binding protein Glutathione peroxidase 1 Fibrinogen Apolipoprotein E	↓ ↑ ↑ ↑
Bozinovski et al. [64]	COPD	Serum	SELDI-TOF MS	Serum amyloid A	↑
Hur et al. [37]	Occupational asthma	BAL	2D-GE Coomassie staining	Ferritin Transferrin	↓ ↑
Lee et al. [33]	COPD	Lung tissue	2D-GE Silver staining	Matrix metalloproteinase 13 Thioredoxin-like 2	↑ ↑
Gomes-Alves et al. [65]	Allergic asthma COPD CF	Serum Nasal epithelial cells	SELDI-TOF MS	Biomarker signatures, one significant identification, hemoglobin subunit beta (↑ in CF as compared to asthma & control)	
Bloemen et al. [58]	Allergic asthma	EBC	LC-MS/MS	Proteolytic peptide profiles discriminate between healthy vs. asthma. No identifications so far	

**Table 2 (Continued)**

Reference	Disease	Sample	Technique	Main findings	Up/down <sup>a</sup>
Gharib et al. [59]	Allergic asthma	Sputum	LC-MS/MS	Calgranulin A	↓
				Calgranulin B	↓
Hu et al. [32]	COPD	Lung tissue	2D-GE	Heat shock protein 27	↑
			MALDI-TOF MS	Cyclophilin A	↑
Ohlmeier et al. [66]	COPD	Sputum	2D-DIGE	Polymeric immunoglobulin receptor	↑
Verrills et al. [34]	COPD	Plasma	2D-DIGE	α2-macroglobulin	↓
	Allergic asthma			haptoglobin	↑
				ceruloplasmin	↓
				hemopexin	↓
Alexandre et al. [56]	COPD	Erythrocyte membrane proteins	O <sup>16</sup> /O <sup>18</sup> labeling	Chorein	↓
			LC-MS/MS	Methemoglobin reductase	↓
Franciosi et al. [31]	COPD	Epithelial lining fluid	iTRAQ	Lactotransferrin	↑
			2D-LC-MS/MS	Cofilin-1	↑
			MALDI-TOF/TOF MS	HMGB1	↓
				Alpha 1-antichymotrypsin	↑
Lee et al. [67]	Uncontrolled asthma	Sputum	2D-GE	S100 calcium binding protein A9	↑
			Coomassie staining		
Merali et al. [68]	COPD	Plasma	1D-GE	Glucose regulated protein 78	↑
			LC-MS/MS	Soluble hemoglobin scavenger receptor	↓
				IL-1 receptor accessory protein	↑
				Macrophage stimulatory protein	↑
Pastor et al. [30]	COPD	BAL	2D-GE	Peroxiredoxin 1	↑
			Sypro Ruby	Heat shock protein 70	↑
				Pyruvate kinase 2	↑
Tu et al. [69]	COPD	BAL	LC-MS/MS	Alcohol dehydrogenase 1B	↑
				Aldehyde dehydrogenase family 3A1	↑
				Aldehyde dehydrogenase (mitochondrial)	↑

<sup>a</sup> Differences presented are always comparisons from a certain disease vs. healthy (e.g. asthma vs. control).

Identification of differentially expressed proteins was performed via mass spectrometry (MALDI-TOF MS or tandem mass spectrometric analysis).

Abbreviations: 2D-GE, two-dimensional gel electrophoresis; 2D-DIGE, two-dimensional difference gel electrophoresis; BAL, bronchoalveolar lavage; CF, cystic fibrosis; COPD, chronic obstructive pulmonary disease; EBC, exhaled breath condensate; LC-MS/MS, liquid chromatography mass spectrometry; PLUNC, palate, lung and nasal epithelium carcinoma associated protein precursor; SELDI-TOF MS, surface enhanced laser desorption/ionization – time of flight mass spectrometry; SSc, systemic sclerosis.

protein biomarkers in blood derived samples (plasma, serum or erythrocytes). These human proteomic studies revealed significant changes in proteins involved in the regulation of inflammation [27–29], oxidative stress [30–32], the acute phase and immune response [31–35] and structural proteins [31,33]. The highlighted proteins in all these studies could be defined as markers of established disease. They are, however, less suited for the detection of early disease onset [16].

The influence of steroid treatment (dexamethasone) on the proteome profile of BAL fluid obtained from murine models, has also been investigated. The level of several proteins, known to be upregulated in allergic asthma, such as chitinase 3-like 3 (Ym1), chitinase 3-like 4 (Ym2) and surfactant protein D were decreased after dexamethasone treatment [36]. Louten et al. [21], using a transgenic mouse model which spontaneously develops asthma, confirmed some of the results. In this study, Ym1 and Ym2 were upregulated both at the protein and mRNA level, comparing control versus asthmatic mice. After dexamethasone treatment, Ym1 mRNA levels decreased whereas Ym2 mRNA levels were not affected by the treatment.

Although most studies have been performed in established allergic asthma (or COPD), a reasonable subset of asthma (9–15%) is attributable to occupational exposures (e.g. animal derived proteins, flour, chemicals, etc.). Only few studies have investigated proteomic changes in occupational asthma. One study by Hur et al. [37] detected significant differences in ferritin and transferrin levels in BAL fluid and serum of diphenylmethane diisocyanate (MDI) asthmatic workers compared to MDI-exposed asymptomatic workers and non-atopic healthy controls. Unfortunately, these results were not confirmed in a more recent study by Sastre et al. [38], investigating workers exposed to toluene diisocyanate (TDI). In one of our studies, we found changes in the levels of ferritin and transferrin in BAL fluid of TDI-asthmatic mice, as shown by Hur et al. The discrepancy between the results obtained by Hur et al. and Sastre et al. concerning biomarkers of diisocyanate asthma are currently a matter of debate since clinical cross-reactivity between diisocyanates (TDI and MDI) is suggested to be common [39]. Moreover, both the study of Hur et al. and Sastre et al. used populations with similar characteristics, although Hur et al. had a considerably larger group of asymptomatic and healthy controls compared to Sastre et al.

Asthma phenotypes share commonalities in their pathology such as oxidative stress, inflammation and non-specific airway hyperreactivity. Nevertheless, some of the proteins such as peroxiredoxins and enolases, are commonly identified regardless of the studied pathology, implying that caution should be taken when attributing the term ‘biomarker’ to a certain protein [40].

So far, proteomic studies conducted in pulmonology have investigated changes in the proteome of healthy subjects versus diseased individuals (asthma, COPD, etc.) using samples from either animal models or patients. Results from a study of our own group in a mouse model of chemical-induced asthma, show that already after two sensitizations, substantial differences occur in the proteome, which we presume relate to the onset of an immune response leading to asthma [41].

### 3. Methodology

Most often 2D-gel electrophoresis (2D-GE) combined with silver or Coomassie staining and a software package, has been used to separate, visualize and analyze differences in protein levels between healthy and diseased subjects (Tables 1 and 2) [26,36,37,42–48]. This is followed by peptide mass fingerprinting or tandem mass spectrometry analysis for the identification of the proteins. The major drawbacks of silver and Coomassie staining are the small linear dynamic range along with the high need for many technical and biological replicates – to correct for the variation between different stained gels – as well as the limited sensitivity of these methods.

The introduction of two-dimensional difference gel electrophoresis (2D-DIGE) has overcome these shortcomings and is now the gold standard among gel-based differential proteomics techniques [49]. This fluorescence-based technique allows the use of multiplexed samples with an internal standard that virtually eliminates gel-to-gel variability, increasing the confidence that uncovered differences are due to real changes, rather than inherent biological variation or experimental variability. Some recent studies in non-malignant lung diseases have applied 2D-DIGE rather than the less preferred silver staining method [20,24,50]. Although 2D-DIGE revolutionized differential analysis in proteomics, this gel-based technique also has some inherent limitations. Particularly, proteins present in extremely low concentrations or proteins that cannot be separated via gel-based techniques due to their physicochemical properties (iso-electric point, hydrophobicity, molecular weight) will not be detected [51,52]. Membrane bound proteins such as surface markers, which could be important in the development of asthma and COPD, are not resolved via gel-based proteomics techniques due to their hydrophobic nature.

Several gel-free proteomics technologies were developed to overcome these limitations. Surface enhanced laser desorption/ionization mass spectrometry (SELDI-TOF) is suited to directly profile subsets of smaller proteins (<20 kDa) based on their physicochemical properties (hydrophobic, metal-binding, etc.) [23,27], however the poor resolution often hampers identification of the proteins.

In other gel-free methods, a protein mixture is enzymatically cleaved into peptides, and subsequently these peptides are separated and identified using liquid chromatography and tandem mass spectrometry (LC-MS/MS) [58,59]. Relative quantitation can be derived by direct comparison of ionic signals produced by peptides in parallel LC-MS analyses. This label-free method uses statistical tools to identify peptides that express consistent differences between samples across multiple runs.

An alternative for relative quantitative analysis is tagging the peptides with chemically almost identical labels that have a mass difference (isotopes). The samples are then combined and analyzed together using LC-MS. The relative abundance of the peptide in the different conditions can be calculated from the ratio of the peak intensity of the different isotopic forms [53]. Isobaric tags for relative and absolute quantification (iTRAQ) is based on covalent labeling of the



N-terminus and side chain amines of peptides from protein digestions with tags of varying mass. Combined with high resolution LC-MS/MS iTRAQ allows the quantitative comparison of protein levels. Franciosi et al. [31] successfully applied this technique in epithelial lining fluid of COPD patients and non-COPD controls and found significantly different levels of lactotransferrin, high-mobility group protein B1, alpha 1-antichymotrypsin and cofilin-1. Stable isotopic labeling by amino acids in cell culture (SILAC) involves the metabolic incorporation of 'normal' or 'heavy' forms ( $C^{13}$  or  $N^{15}$ ) of arginine or lysine, supplied in culture medium, in newly synthesized proteins. The normal and heavy forms of the peptides appear as two distinct peaks in MS and relative quantitation is achieved by comparing the signal intensities. SILAC has been successfully used in studies focusing on lung cancer [54,55]. Alexandre et al. [56], used differential oxygen ( $O^{16}/O^{18}$ ) labeling to quantify differences in erythrocyte membrane proteins of COPD patients compared to healthy controls. Differential  $O^{16}/O^{18}$  labeling occurs at the C-terminal carboxyl group of peptides, derived from the proteolytic digestion of proteins, where two  $O^{16}$  atoms are replaced by  $O^{18}$  atoms. The resulting mass shift permits the relative quantitation of proteins between two conditions [57].

These gel-free methods also have their inherent limitations. A typical tissue extract contains a few thousands of proteins that after digestion will result in a more complex peptides mixture, while the number of peptides that can be identified due to co-elution and ion suppression is limited [58]. Posttranslational modifications such as phosphorylations, often important in the regulation of protein function, will be visualized on 2D gels by different spots as a result of shift in isoelectric point (pI). In gel-free proteomics, pI information, together with protein molecular weight information, is lost.

#### 4. Mice vs. men

One can debate if human samples are preferable over samples obtained from mouse models. Nonetheless, significant understanding on the pathogenesis of asthma and COPD has been derived from animal experiments, particularly in mice and rats [25,26,70]. Several mouse specific immunological tools and the availability to use transgenic mice (e.g. TSLSP Tg mice) favor this species [21]. Moreover, 99% of mouse genes have homologues in humans, facilitating proteomics approaches [71]. Also, ethical and moral issues prevent profound mechanistic investigations in humans [72]. Animals are generally kept in clean animal houses, limiting the environmental factors to which they are exposed and facilitate the exposure to exact concentrations of a known substance (e.g. allergen or tobacco smoke) [73]. In animals, invasive samples, such as lung tissue, lymphoid tissue and BAL fluid are more easily obtained than in humans.

Unfortunately, the results obtained from animal experiments cannot directly be extrapolated to humans since important differences in airway morphology and development exist between mice and humans. Moreover, the initiation, development and responses of the innate and adaptive immune system differ substantially between mice and men

[74]. No mouse model is currently available that matches all phenotypic characteristics of human asthma [75–77], nor COPD [78]. Most mouse models are an acute model of asthma, lacking chronicity and airway remodeling, two characteristics of human asthma. The choice of mouse strain has also been demonstrated to have an influence on the physiological parameters (cytokines, neutrophils, eosinophils, etc.) [79–81]. Yet, it is possible to induce several characteristics of the disease such as lymphocyte activation and IgE production, along with airway hyperresponsiveness, airway inflammation and in some studies airway remodeling [24,82–87].

Bearing in mind these shortcomings, animal models can be seen as a platform to generate and test hypotheses which need validation in humans [88]. Despite the significant results obtained from animal studies, translation to human disease is difficult and time consuming. Hackam et al. [11] have reviewed highly cited animal studies, showing that only one out of three animal studies evaluating medical interventions for human diseases have been replicated in clinical trials.

#### 5. Sample origin

Another important issue concerns the best choice of tissue for analysis. So far, lung tissue has been the most used sample in murine models of asthma and COPD for obvious reasons [23,24,26,42,43,45,89]. In animals, it is possible to obtain samples requiring more invasive techniques and, moreover, lung tissue will possibly contain high concentrations of potential biomarkers specific for lung diseases. Nevertheless, studying proteome changes in lung tissue by gel-based techniques is limited to the soluble proteins. It is important to validate the biomarkers obtained in lung tissue, using more accessible samples, such as BAL fluid and to see how well data of both compartments are correlated [90]. BAL fluid provides the most faithful reflection of the protein composition of the pulmonary lumen and is therefore the preferred sample in humans [91–93]. However, proteomic analysis of BAL fluid is complicated by a low protein abundance and high salt content and possible contamination by serum proteins, as a result from leakage during lung inflammation [94]. Although serum proteins in BAL could also reflect the current pulmonary situation, albumin and immunoglobulin depletion strategies have been developed and applied to improve the quality of proteomic studies [20].

Recently, in humans, the use of less invasive techniques such as induced sputum and exhaled breath condensate (EBC) have also been evaluated [27,58]. Induced sputum and EBC can be routinely collected from patients during follow-up, but the use of these samples in proteomics studies is scarce due to the very low protein content and needs to be optimized.

Strikingly, only few of the studies listed in Tables 1 and 2 used plasma or serum samples [20,41,63,64]. Serum is an ideal biological sample that contains an archive of information due to the presence of a variety of proteins released by diseased tissue [95]. Nevertheless, proteomic analysis of serum has several advantages and disadvantages. Serum is easy to acquire, can be obtained at different time points (e.g. during follow-up) and has a high protein content. However, high-abundant proteins (albumin, IgG and transferrin) can possibly mask potential



biomarkers with a relative low abundance, and serum is also highly variable [96].

It seems that the samples with the highest protein content show fewer significant differential proteins, giving the impression that the complexity reduces the sensitivity. Therefore, enrichments of subsets of cells, such as T- and B-lymphocytes, or macrophages and leucocytes, could reduce the complexity and facilitate proteomic analysis. On the other hand, a lot of starting material (*cfr.* mice) is required to have enough cells to perform these experiments, which is a major limiting factor. In humans, this can be overcome because a large amount of blood can be sampled to isolate specific lymphocyte subsets, as done by Jeong et al. [47], who compared human peripheral T-lymphocytes from asthma patients and healthy controls.

## 6. Interpretation of results

Most often, the end-result of a proteomic study is the identification of ten to hundreds of differentially expressed proteins which are presented in a table. A few of these proteins are selected based on literature and biological relevance. Several proteins, such as annexins, peroxiredoxins and haptoglobins, are differentially expressed in different studies when comparing all the results of the papers described in Tables 1 and 2. One should question if these proteins could be called biomarkers since these proteins are repeatedly identified throughout different research domains and are therefore called ‘d  ja-vu’-proteins [40]. Chitinase 3-like 3 (Ym1) and Chitinase 3-like 4 (Ym2) are both members of the chitinase family, responsible for the degradation of chitin which is present in the exoskeleton of arthropods and in cell membranes of fungi. Although chitin is not present in mammals they are regularly associated with asthma in murine models [21,45,60]. Although no human orthologues have been identified for Ym1 and Ym2, other human chitinases have been demonstrated to be involved in the pathogenesis of allergic disease and COPD.

In summary, drawing general conclusions from common identified proteins is challenging, because of the variety of mouse strains, tissues and treatment protocols (difference in timing, dosage and duration) that are used to induce asthma or COPD (e.g. acute versus chronic). In addition, substantial differences in physiology between mice and man set hurdles in extrapolating the results obtained from murine models to humans.

## 7. Future

The rationale for choosing proteomics approaches, is to discover novel target proteins as biomarkers for disease identification, as well as possible new markers with the focus on refining treatment. Although different proteins are highlighted in Tables 1 and 2 that could be linked with the phenotypical characteristics of asthma or COPD in murine models and humans, functional biological validation – specifically for murine studies – is nearly always lacking, thereby indicating the long way that is still to go in proteomics biomarker research. One exception is the 2D-DIGE study of lung tissue of Calvo et al. [24], who discovered the involvement of a 78 kDa glucose-regulated protein precursor (Grp78)

in OVA-induced asthma in mice. They silenced down the gene responsible for this protein using anti-Grp78 siRNA treatment in OVA-sensitized and challenged mice, resulting in decreased airway hyperreactivity (AHR) and eosinophilic inflammation. This type of validation is valuable in determining the biological relevance of the studied proteins as possible therapeutic targets. On the other hand, proteomics studies are often used to screen for markers of disease. Clinically useful biomarkers would provide earlier and better diagnosis of patients, permitting treatments to be initiated in early stages of the disease, when the chance of success is greatest [96,97]. Several proteins such as ferritin, transferrin and calcyclin were proposed as general markers of lung disease [23,37]. As already indicated, Hur et al. described ferritin and transferrin as possible biomarkers of MDI-induced asthma, but Sastre et al. could not verify these markers. Moreover, ferritin and transferrin are involved in a broad range of biological processes, limiting their use as specific biomarkers. More efforts should be focused to find disease-specific, unique biomarkers for lung diseases. Perhaps future efforts should be less dedicated to known proteins found differentially in these studies, but more to explore the proteins with unknown ontology. This could lead to really novel and innovating biomarkers and perhaps also therapeutic targets.

Furthermore, current state-of-the-art diagnosis and management schemes of asthma are suboptimal since the incidence of asthma has risen by 250% over the last two decades [16]. There is an urgent need for early diagnosis since asthma generally starts early during childhood. Occupational asthma often is exclusively related to exposures on the work floor and thereby only develops at a later stage of life. However, the onset of allergic OA is always preceded by a complaints-free latency period in which workers are getting sensitized, as a result of complex immunological and molecular processes that are not identified by currently used diagnostic tools but which could be picked up by ‘-omics’ techniques.

In addition, peptide hormones, neuropeptides, cytokines and chemokines are involved in the regulation of numerous physiological processes and the onset, maintenance and progression of diseases such as asthma. These peptides are generally not detected by proteomics techniques either due to their size (<20 kDa) or to their low abundance and therefore are a yet unexplored pool of possible new biomarkers [98]. Peptidomics techniques, using a combination of liquid chromatography and mass spectrometry can be used to identify peptides. Nonetheless, quantitative peptidomics is still a major challenge.

## 8. Conclusion

There is an increasing trend in the use of proteomics to explore new biomarkers of asthma or COPD. Although different murine models have been developed to study these lung diseases, and proteomics studies have been performed, functional validation of the identified candidate biomarkers or translation into clinical studies is often lacking. Shifting the focus from disease toward the early events during disease manifestation (e.g. sensitization) and the investigation of peptides as a yet unexplored pool of interesting biomarkers

would bear great potential in the identification of possible new, clinically useful biomarkers.

### Conflict of interest statement

The authors declare that they have no conflicts of interest.

### Transparency document

The [Transparency document](#) associated with this article can be found in the online version.

### REFERENCES

- [1] Ahram M, Petricoin EF. Proteomics discovery of disease biomarkers. *Biomark Insights* 2008;3:325–33.
- [2] Bogaerts A, Temmerman L, Boerjan B, Husson SJ, Schoofs L, Verleyen P. A differential proteomics study of *Caenorhabditis elegans* infected with *Aeromonas hydrophila*. *Dev Comp Immunol* 2010;34(6):690–8.
- [3] Griffith OL, Gray JW. Omic approaches to preventing or managing metastatic breast cancer. *Breast Cancer Res* 2011;13(6):230.
- [4] Sun C, Rosendahl AH, Ansari D, Andersson R. Proteome-based biomarkers in pancreatic cancer. *World J Gastroenterol* 2011;17(44):4845–52.
- [5] Goh WW, Lee YH, Chung M, Wong L. How advancement in biological network analysis methods empowers proteomics. *Proteomics* 2012;12(4–5):550–63.
- [6] Decramer M, Janssens W, Miravittles M. Chronic obstructive pulmonary disease. *Lancet* 2012;379(9823):1341–51.
- [7] Cottini M, Asero R. Asthma phenotypes today. *Eur Ann Allergy Clin Immunol* 2013;45(1):17–24.
- [8] Martinez FD, Vercelli D. Asthma. *Lancet* 2013;382(9901):1360–72.
- [9] Vahakangas K. Research ethics in the post-genomic era. *Environ Mol Mutagen* 2013;54(7):599–610.
- [10] Hirsch J, Hansen KC, Burlingame AL, Matthay MA. Proteomics: current techniques and potential applications to lung disease. *Am J Physiol Lung Cell Mol Physiol* 2004;287(1):L1–23.
- [11] Hackam DG, Redelmeier DA. Translation of research evidence from animals to humans. *J Am Med Assoc* 2006;296(14):1731–2.
- [12] Maddox L, Schwartz DA. The pathophysiology of asthma. *Annu Rev Med* 2002;53:477–98.
- [13] Busse WW, Rosenwasser LJ. Mechanisms of asthma. *J Allergy Clin Immunol* 2003;111:799–804.
- [14] Lemanske Jr RF, Busse WW. 6. Asthma. *J Allergy Clin Immunol* 2003;111(2 Suppl.):S502–19.
- [15] Svanes C, Sunyer J, Plana E, Dharmage S, Heinrich J, Jarvis D, et al. Early life origins of chronic obstructive pulmonary disease. *Thorax* 2010;65(1):14–20.
- [16] Lin JL, Bonnicksen MH, Nogeh EU, Raftery MJ, Thomas PS. Proteomics in detection and monitoring of asthma and smoking-related lung diseases. *Expert Rev Proteomics* 2010;7(3):361–72.
- [17] Wolthers OD. Eosinophil granule proteins in the assessment of airway inflammation in pediatric bronchial asthma. *Pediatr Allergy Immunol* 2003;14(4):248–54.
- [18] Hopkins AL, Groom CR. The druggable genome. *Nat Rev Drug Discov* 2002;1(9):727–30.
- [19] Signor L, Tigani B, Beckmann N, Falchetto R, Stoeckli M. Two-dimensional electrophoresis protein profiling and identification in rat bronchoalveolar lavage fluid following allergen and endotoxin challenge. *Proteomics* 2004;4(7):2101–10.
- [20] Haenen S, Vanoirbeek JA, De Vooght V, Maes E, Schoofs L, Nemery B, et al. Proteome analysis of multiple compartments in a mouse model of chemical-induced asthma. *J Proteome Res* 2010;9(11):5868–76.
- [21] Louten J, Mattson JD, Malinao MC, Li Y, Emson C, Vega F, et al. Biomarkers of disease and treatment in murine and cynomolgus models of chronic asthma. *Biomark Insights* 2012;7:87–104.
- [22] Wu J, Kobayashi M, Sousa EA, Liu W, Cai J, Goldman SJ, et al. Differential proteomic analysis of bronchoalveolar lavage fluid in asthmatics following segmental antigen challenge. *Mol Cell Proteomics* 2005;4(9):1251–64.
- [23] Calvo FQ, Fillet M, de Seny D, Meuwis MA, Maree R, Crahay C, et al. Biomarker discovery in asthma-related inflammation and remodeling. *Proteomics* 2009;9(8):2163–70.
- [24] Calvo FQ, Fillet M, Renaut J, Crahay C, Gueders M, Hacha J, et al. Potential therapeutic target discovery by 2D-DIGE proteomic analysis in mouse models of asthma. *J Proteome Res* 2011;10(9):4291–301.
- [25] Zhang S, Xu N, Nie J, Dong L, Li J, Tong J. Proteomic alteration in lung tissue of rats exposed to cigarette smoke. *Toxicol Lett* 2008;178(3):191–6.
- [26] Xu NY, Zhang SP, Dong L, Nie JH, Tong J. Proteomic analysis of lung tissue of rats exposed to cigarette smoke and radon. *J Toxicol Environ Health A* 2009;72(11–12):752–8.
- [27] Gray RD, MacGregor G, Noble D, Imrie M, Dewar M, Boyd AC, et al. Sputum proteomics in inflammatory and suppurative respiratory diseases. *Am J Respir Crit Care Med* 2008;178(5):444–52.
- [28] Merkel D, Rist W, Seither P, Weith A, Lenter MC. Proteomic study of human bronchoalveolar lavage fluids from smokers with chronic obstructive pulmonary disease by combining surface-enhanced laser desorption/ionization-mass spectrometry profiling with mass spectrometric protein identification. *Proteomics* 2005;5(11):2972–80.
- [29] Steiling K, Kadar AY, Bergerat A, Flanigan J, Sridhar S, Shah V, et al. Comparison of proteomic and transcriptomic profiles in the bronchial airway epithelium of current and never smokers. *PLoS One* 2009;4(4):e5043.
- [30] Pastor MD, Nogal A, Molina-Pinelo S, Melendez R, Romero-Romero B, Mediano MD, et al. Identification of oxidative stress related proteins as biomarkers for lung cancer and chronic obstructive pulmonary disease in bronchoalveolar lavage. *Int J Mol Sci* 2013;14(2):3440–55.
- [31] Franciosi L, Govorukhina N, Fusetti F, Poolman B, Lodewijk ME, Timens W, et al. Proteomic analysis of human epithelial lining fluid by microfluidics-based nanoLC-MS/MS: a feasibility study. *Electrophoresis* 2013;34(18):2683–94.
- [32] Hu R, Ouyang Q, Dai A, Tan S, Xiao Z, Tang C. Heat shock protein 27 and cyclophilin A associate with the pathogenesis of COPD. *Respirology* 2011;16(6):983–93.
- [33] Lee EJ, In KH, Kim JH, Lee SY, Shin C, Shim JJ, et al. Proteomic analysis in lung tissue of smokers and COPD patients. *Chest* 2009;135(2):344–52.
- [34] Verrills NM, Irwin JA, He XY, Wood LG, Powell H, Simpson JL, et al. Identification of novel diagnostic biomarkers for asthma and chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2011;183(12):1633–43.
- [35] Tewari AK, Popova-Butler A, El-Mahdy MA, Zweier JL. Identification of differentially expressed proteins in blood plasma of control and cigarette smoke-exposed mice by 2-D DIGE/MS. *Proteomics* 2011;11(10):2051–62.
- [36] Zhao J, Yeong LH, Wong WS. Dexamethasone alters bronchoalveolar lavage fluid proteome in a mouse asthma model. *Int Arch Allergy Immunol* 2007;142(3):219–29.

- [37] Hur GY, Choi GS, Sheen SS, Lee HY, Park HJ, Choi SJ, et al. Serum ferritin and transferrin levels as serologic markers of methylene diphenyl diisocyanate-induced occupational asthma. *J Allergy Clin Immunol* 2008;122(4):774–80.
- [38] Sastre J, Sastre B, Fernandez-Nieto M, Perez-Camo I, Sanchez JJ, Del Pozo V. Serum ferritin and transferrin levels are not serologic markers of toluene diisocyanate-induced occupational asthma. *J Allergy Clin Immunol* 2010;125(3):762–4.
- [39] Vogelmeier C, Baur X, Fruhmman G. Isocyanate-induced asthma: results of inhalation tests with TDI, MDI and methacholine. *Int Arch Occup Environ Health* 1991;63(1):9–13.
- [40] Petrak J, Ivanek R, Toman O, Cmejla R, Cmejlova J, Vyoral D, et al. Deja vu in proteomics. A hit parade of repeatedly identified differentially expressed proteins. *Proteomics* 2008;8(9):1744–9.
- [41] Haenen S, Clynen E, De Vooght V, Schoofs L, Nemery B, Hoet PH, et al. Proteome changes in auricular lymph nodes and serum after dermal sensitization to toluene diisocyanate in mice. *Proteomics* 2012;12(23–24):3548–58.
- [42] Houtman R, Krijgsveld J, Kool M, Romijn EP, Redegeld FA, Nijkamp FP, et al. Lung proteome alterations in a mouse model for nonallergic asthma. *Proteomics* 2003;3(10):2008–18.
- [43] Roh GS, Shin Y, Seo SW, Yoon BR, Yeo S, Park SJ, et al. Proteome analysis of differential protein expression allergen-induced asthmatic mice lung after dexamethasone treatment. *Proteomics* 2004;4(11):3318–27.
- [44] Candiano G, Bruschi M, Pedemonte N, Caci E, Liberatori S, Bini L, et al. Gelsolin secretion in interleukin-4-treated bronchial epithelia and in asthmatic airways. *Am J Respir Crit Care Med* 2005;172(9):1090–6.
- [45] Jeong H, Rhim T, Ahn MH, Yoon PO, Kim SH, Chung IY, et al. Proteomic analysis of differently expressed proteins in a mouse model for allergic asthma. *J Korean Med Sci* 2005;20(4):579–85.
- [46] Larsen K, Malmstrom J, Wildt M, Dahlqvist C, Hansson L, Marko-Varga G, et al. Functional and phenotypical comparison of myofibroblasts derived from biopsies and bronchoalveolar lavage in mild asthma and scleroderma. *Respir Res* 2006;7:11.
- [47] Jeong HC, Lee SY, Lee EJ, Jung KH, Kang EH, Lee SY, et al. Proteomic analysis of peripheral T-lymphocytes in patients with asthma. *Chest* 2007;132(2):489–96.
- [48] Ohlmeier S, Vuolanto M, Toljamo T, Vuopala K, Salmenkivi K, Myllarniemi M, et al. Proteomics of human lung tissue identifies surfactant protein A as a marker of chronic obstructive pulmonary disease. *J Proteome Res* 2008;7(12):5125–32.
- [49] Rozanas CR, Loyland SM. Capabilities using 2-D DIGE in proteomics research: the new gold standard for 2-D gel electrophoresis. *Methods Mol Biol* 2008;441:1–18.
- [50] Greenlee KJ, Corry DB, Engler DA, Matsunami RK, Tessier P, Cook RG, et al. Proteomic identification of in vivo substrates for matrix metalloproteinases 2 and 9 reveals a mechanism for resolution of inflammation. *J Immunol* 2006;177(10):7312–21.
- [51] Baggerman G, Vierstraete E, De Loof A, Schoofs L. Gel-based versus gel-free proteomics: a review. *Comb Chem High Throughput Screen* 2005;8(8):669–77.
- [52] Gygi SP, Corthals GL, Zhang Y, Rochon Y, Aebersold R. Evaluation of two-dimensional gel electrophoresis-based proteome analysis technology. *Proc Natl Acad Sci U S A* 2000;97(17):9390–5.
- [53] Pastor MD, Nogal A, Molina-Pinelo S, Carnero A, Paz-Ares L. Proteomic biomarkers in lung cancer. *Clin Transl Oncol* 2013;15(9):671–82.
- [54] Schweppe DK, Rigas JR, Gerber SA. Quantitative phosphoproteomic profiling of human non-small cell lung cancer tumors. *J Proteomics* 2013;91:286–96.
- [55] Zhang W, Wei Y, Ignatchenko V, Li L, Sakashita S, Pham NA, et al. Proteomic profiles of human lung adeno and squamous cell carcinoma using super-SILAC and label-free quantification approaches. *Proteomics* 2014;14(6):795–803.
- [56] Alexandre BM, Charro N, Blonder J, Lopes C, Azevedo P, Bugalho de Almeida A, et al. Profiling the erythrocyte membrane proteome isolated from patients diagnosed with chronic obstructive pulmonary disease. *J Proteomics* 2012;76(Spec No.):259–69.
- [57] Ye X, Luke B, Andresson T, Blonder J. <sup>18</sup>O stable isotope labeling in MS-based proteomics. *Brief Funct Genomic Proteomic* 2009;8(2):136–44.
- [58] Bloemen K, Van Den Heuvel R, Govarts E, Hooyberghs J, Nelen V, Witters E, et al. A new approach to study exhaled proteins as potential biomarkers for asthma. *Clin Exp Allergy* 2011;41(3):346–56.
- [59] Gharib SA, Nguyen EV, Lai Y, Plampin JD, Goodlett DR, Hallstrand TS. Induced sputum proteome in healthy subjects and asthmatic patients. *J Allergy Clin Immunol* 2011;128(6):1176–84.
- [60] Zhao J, Zhu H, Wong CH, Leung KY, Wong WS. Increased lungkine and chitinase levels in allergic airway inflammation: a proteomics approach. *Proteomics* 2005;5(11):2799–807.
- [61] Liu H, Zhou LF, Qian Z, Qian FH, Yin KS, Mao H, et al. Increased RhoGDI(2) and peroxiredoxin 5 levels in asthmatic murine model of beta(2)-adrenoceptor desensitization: a proteomics approach. *Chin Med J* 2008;121(4):355–62.
- [62] Larsen K, Macleod D, Nihlberg K, Gurcan E, Bjermer L, Marko-Varga G, et al. Specific haptoglobin expression in bronchoalveolar lavage during differentiation of circulating fibroblast progenitor cells in mild asthma. *J Proteome Res* 2006;5(6):1479–83.
- [63] Bandow JE, Baker JD, Berth M, Painter C, Sepulveda OJ, Clark KA, et al. Improved image analysis workflow for 2-D gels enables large-scale 2-D gel-based proteomics studies – COPD biomarker discovery study. *Proteomics* 2008;8(15):3030–41.
- [64] Bozinovski S, Hutchinson A, Thompson M, Macgregor L, Black J, Giannakis E, et al. Serum amyloid a is a biomarker of acute exacerbations of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2008;177(3):269–78.
- [65] Gomes-Alves P, Imrie M, Gray RD, Nogueira P, Giordis S, Pacheco P, et al. SELDI-TOF biomarker signatures for cystic fibrosis, asthma and chronic obstructive pulmonary disease. *Clin Biochem* 2010;43(1–2):168–77.
- [66] Ohlmeier S, Mazur W, Linja-Aho A, Louhelainen N, Ronty M, Toljamo T, et al. Sputum proteomics identifies elevated PIGR levels in smokers and mild-to-moderate COPD. *J Proteome Res* 2012;11(2):599–608.
- [67] Lee TH, Jang AS, Park JS, Kim TH, Choi YS, Shin HR, et al. Elevation of S100 calcium binding protein A9 in sputum of neutrophilic inflammation in severe uncontrolled asthma. *Ann Allergy Asthma Immunol* 2013;111(4):268–75.
- [68] Merali S, Barrero CA, Bowler RP, Chen DE, Criner G, Braverman A, et al. Analysis of the plasma proteome in COPD: novel low abundance proteins reflect the severity of lung remodeling. *COPD* 2014;11(2):177–89.
- [69] Tu C, Mammen MJ, Li J, Shen X, Jiang X, Hu Q, et al. Large-scale, ion-current-based proteomics investigation of bronchoalveolar lavage fluid in chronic obstructive pulmonary disease patients. *J Proteome Res* 2014;13(2):627–39.
- [70] Irvin CG. Using the mouse to model asthma: the cup is half full and then some. *Clin Exp Allergy* 2008;38(5):701–3.

- [71] Waterston RH, Lindblad-Toh K, Birney E, Rogers J, Abril JF, Agarwal P, et al. Initial sequencing and comparative analysis of the mouse genome. *Nature* 2002;420(6915):520–62.
- [72] Johnson VJ, Matheson JM, Luster MI. Animal models for diisocyanate asthma: answers for lingering questions. *Curr Opin Allergy Clin Immunol* 2004;4:105–10.
- [73] Wright JL, Churg A. Animal models of COPD: barriers, successes, and challenges. *Pulm Pharmacol Ther* 2008;21(5):696–8.
- [74] Mestas J, Hughes CC. Of mice and not men: differences between mouse and human immunology. *J Immunol* 2004;172(5):2731–8.
- [75] Kips JC, Anderson GP, Fredberg JJ, Herz U, Inman MD, Jordana M, et al. Murine models of asthma. *Eur Respir J* 2003;22(2):374–82.
- [76] Shapiro SD. Animal models of asthma: Pro: allergic avoidance of animal (model[s]) is not an option. *Am J Respir Crit Care Med* 2006;174(11):1171–3.
- [77] Ware LB. Modeling human lung disease in animals. *Am J Physiol Lung Cell Mol Physiol* 2008;294(2):L149–50.
- [78] Brusselle GG, Bracke KR, Maes T, D'hulst AI, Moerloose KB, Joos GF, et al. Murine models of COPD. *Pulm Pharmacol Ther* 2006;19(3):155–65.
- [79] De Vooght V, Vanoirbeek JA, Luyts K, Haenen S, Nemery B, Hoet PH. Choice of mouse strain influences the outcome in a mouse model of chemical-induced asthma. *PLoS One* 2010;5(9):e12581.
- [80] Rueff-Barroso CR, Trajano ET, Alves JN, Paiva RO, Lanzetti M, Pires KM, et al. Organ-related cigarette smoke-induced oxidative stress is strain-dependent. *Med Sci Monit* 2010;16(7):BR218–26.
- [81] Vecchio D, Arezzini B, Pecorelli A, Valacchi G, Martorana PA, Gardi C. Reactivity of mouse alveolar macrophages to cigarette smoke is strain dependent. *Am J Physiol Lung Cell Mol Physiol* 2010;298(5):L704–13.
- [82] Herrick CA, Xu L, Wisnewski AV, Das J, Redlich CA, Bottomly K. A novel mouse model of diisocyanate-induced asthma showing allergic-type inflammation in the lung after inhaled antigen challenge. *J Allergy Clin Immunol* 2002;109(5):873–8.
- [83] Vanoirbeek JAJ, Tarkowski M, Ceuppens JL, Verbeken EK, Nemery B, Hoet PHM. Respiratory response to toluene diisocyanate depends on prior frequency and concentration of dermal sensitization in mice. *Toxicol Sci* 2004;80(2):310–21.
- [84] Matheson JM, Johnson VJ, Vallyathan V, Luster MI. Exposure and immunological determinants in a murine model for toluene diisocyanate (TDI) asthma. *Toxicol Sci* 2005;84(1):88–98.
- [85] Tarkowski M, Vanoirbeek JA, Vanhooren HM, De Vooght V, Mercier CM, Ceuppens J, et al. Immunological determinants of ventilatory changes induced in mice by dermal sensitization and respiratory challenge with toluene diisocyanate. *Am J Physiol Lung Cell Mol Physiol* 2007;292(1):L207–14.
- [86] De Vooght V, Vanoirbeek JA, Haenen S, Verbeken E, Nemery B, Hoet PH. Oropharyngeal aspiration: an alternative route for challenging in a mouse model of chemical-induced asthma. *Toxicology* 2009;259(1–2):84–9.
- [87] Vanoirbeek JA, Tarkowski M, De Vooght V, Nemery B, Hoet PH. Immunological determinants in a mouse model of chemical-induced asthma after multiple exposures. *Scand J Immunol* 2009;70(1):25–33.
- [88] Wenzel S, Holgate ST. The mouse trap: it still yields few answers in asthma. *Am J Respir Crit Care Med* 2006;174(11):1173–6.
- [89] Zhang SP, Xu NY, Nie JH, Dong L, Li JX, Tong J. Proteomic alteration in lung tissue of rats exposed to cigarette smoke. *Toxicol Lett* 2008;178(3):191–6.
- [90] Zhang L, Wang M, Kang X, Boontheung P, Li N, Nel AE, et al. Oxidative stress and asthma: proteome analysis of chitinase-like proteins and FIZZ1 in lung tissue and bronchoalveolar lavage fluid. *J Proteome Res* 2009;8(4):1631–8.
- [91] Noel-Georis I, Bernard A, Falmagne P, Wattiez R. Database of bronchoalveolar lavage fluid proteins. *J Chromatogr B Analyt Technol Biomed Life Sci* 2002;771(1–2):221–36.
- [92] Magi B, Bargagli E, Bini L, Rottoli P. Proteome analysis of bronchoalveolar lavage in lung diseases. *Proteomics* 2006;6(23):6354–69.
- [93] Hu S, Loo JA, Wong DT. Human body fluid proteome analysis. *Proteomics* 2006;6(23):6326–53.
- [94] Ratjen F, Havers W, Braun J. Intrapulmonary protein leakage in immunocompromised children and adults with pneumonia. *Thorax* 1999;54(5):432–6.
- [95] Ray S, Reddy PJ, Jain R, Gollapalli K, Moiyadi A, Srivastava S. Proteomic technologies for the identification of disease biomarkers in serum: advances and challenges ahead. *Proteomics* 2011;11(11):2139–61.
- [96] Blonder J, Issaq HJ, Veenstra TD. Proteomic biomarker discovery: it's more than just mass spectrometry. *Electrophoresis* 2011;32(13):1541–8.
- [97] Schilsky RL. Personalized medicine in oncology: the future is now. *Nat Rev Drug Discov* 2010;9(5):363–6.
- [98] Clynen E, Baggerman G, Husson SJ, Landuyt B, Schoofs L. Peptidomics in drug research. *Expert Opin Drug Discov* 2008;3(4):425–40.